#### § 180.655

isopyrazam, 3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxylic acid (9-isopropyl-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl)-amide, in or on the following commodity.

Commodity	Parts per million
Banana <sup>1</sup>	0.05

- <sup>1</sup>There is no U.S. registration for use of isopyrazam on hanana
- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[76 FR 61596, Oct. 5, 2011]

### § 180.655 Flazasulfuron; tolerances for residues.

(a) General. Tolerances are established for residues of flazasulfuron, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only flazasulfuron (N-[[(4,6-dimethoxy-2-

pyrimidinyl)amino]carbonyl]-3-(trifluoromethyl)-2pyridinesulfonamide).

Commodity	Parts per million
Fruit, citrus, group 10-10	0.01
Grape	0.01
Sugarcane	0.01

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[77 FR 10968, Feb. 24, 2012]

# § 180.656 Amisulbrom; tolerances for residues.

(a) General. Tolerances are established for residues of the fungicide amisulbrom, including its metabolites and degradates, in or on the commodities listed below. Compliance with the tolerance levels is to be determined by measuring only amisulbrom, 3-[(3-bromo-6-fluoro-2-methyl-1H-indole-1-yl) sulfonyl]-N, N-dimethyl-1H-1, 2, 4-triazole-1-sulfonamide].

Commodity <sup>1</sup>	Parts per million
Grape	0.40
Grape, raisin	1.0
Tomato	0.50
Tomato, paste	1.2

- $^{\rm 1} There$  is no U.S. registration for use of amisulbrom on grape or tomato.
- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[76 FR 59914, Sept. 28, 2011]

## § 180.657 Metaflumizone; tolerances for residues.

(a) General. Tolerances are established for residues of the insecticide metaflumizone, including its metabolites and degradates, in or on the commodities listed in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only the sum of metaflumizone (E and Z isomers; 2-[2-(4-cyanophenyl)-1-[3-

(trifluoromethyl) phenyl]ethylidene]-N-[4-(trifluoromethoxy)phenyl]

hydrazinecarboxamide) and its metabolite 4-{2-oxo-2-[3-(trifluoromethyl) phenyl]ethyl}-benzonitrile, calculated as the stoichiometric equivalent of metaflumizone, in or on the following commodities:

Commodity	Parts per million
Almond, hulls Fruit, citrus, group 10 Grape	0.04 0.04 0.04
Nut, tree, group 14	0.04

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[77 FR 10386, Feb. 22, 2012]

# §180.658 Penthiopyrad; tolerances for residues.

(a) General. (1) Tolerances are established for residues of penthiopyrad, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance

### **Environmental Protection Agency**

levels specified below is to be determined by measuring only penthiopyrad (N-[2-(1,3-dimethylbutyl)-3-thienyl]-1methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide).

•	
Commodity	Parts per million
Alfalfa, forage	7.0
Alfalfa, hay	20
Almond, hulls	6.0
Apple, wet pomace	1.5
Barley, grain	0.15
Barley, hay	80
Barley, milled byproducts Barley, straw	0.90 1.0
Beet, sugar, dried pulp	1.5
Beet, sugar, roots	0.5
Berry, low growing, subgroup 13-07G	3.0
Brassica, head and stem, subgroup 5A	5.0
Brassica, leafy greens, subgroup 5B	50
Buckwheat, grain	0.15
Canola	1.5
Corn, field, grain	40 0.01
Corn, field, grain  Corn, field, refined oil	0.01
Corn field stover	15
Corn, field, stover	0.01
Corn, sweet, kernel plus cob with husks re-	
moved	0.01
Cotton, seed	1.5
Cotton, gin byproducts	15
Fruit, pome, group 11–10	0.50
Fruit, stone, group 12	4.0
Grain, aspirated fractions	30
Nut, tree, group 14	0.80 0.06
Oat, forage	40
Oat, grain	0.15
Oat, hay	80
Oat. straw	1.0
Pea and bean, dried shelled, except soybean,	
subgroup 6C	0.40
Peanut	0.04
Peanut, hay	30
Peanut, refined oil	0.06 0.06
Potato, processed potato waste	0.00
Rye, forage	40
Rve. grain	0.15
Rye, grain	1.0
Sorghum, forage	40
Sorghum, grain, grain	0.80
Sorghum, stover	15
Soybean, seed	0.40
Sunflower, seed	1.5
Teosinte, grain Tomato, paste	0.15 3.5
Triticale, forage	40
Triticale, rorage	0.15
Triticale, grain Triticale, hay	80
Triticale, straw	1.0
Vegetable, bulb, group 3-07	3.0
Vegetable, cucurbit, group 9	0.60
Vegetable, foliage of legume, group 7, hay	200
Vegetable, foliage of legume, group 7, vines/for-	
age	50
Vegetable, fruiting, group 8–10	3.0
Vegetable, leafy, except brassica, group 4	30 50
Vegetable, leaves of root and tuber, group 2	50 4.0
Vegetable, legume, edible podded, subgroup 6A Vegetable, legume, succulent shelled, subgroup	4.0
6B	0.40
Vegetable, root, subgroup 1B, except sugar	JFU
beet	3.0

Commodity	Parts per million
Vegetable, tuber and corm, subgroup 1C	0.06 40 0.15 80 0.30 1.0

(2) Tolerances are established for residues of penthiopyrad, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum penthiopyrad (N-[2-(1,3-dimethylbutyl)-3-thienyl]-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4carboxamide) and its metabolite (1methyl-3-trifluoromethyl-1H-pyrazole-4-carboxamide), calculated as the stoichiometric equivalent of penthiopyrad, in or on the commodity.

Commodity	Parts per million
Cattle, fat	0.03
Cattle, meat	0.03
Cattle, meat byproducts	0.09
Goat, fat	0.03
Goat, meat	0.03
Goat, meat byproducts	0.09
Horse, fat	0.03
Horse, meat	0.03
Horse, meat byproducts	0.09
Milk	0.02
Sheep, fat	0.03
Sheep, meat	0.03
Sheep, meat byproducts	0.09

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[77 FR 14297, Mar. 9, 2012]

#### §180.659 Pyroxasulfone; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the herbicide pyroxasulfone, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of pyroxasulfone, 3-[[[5-(difluoromethoxy)-1-methyl-3-(trifluoromethyl)-1*H*-pyrazol-4-yl] methyl]sulfonyl]-4,5-dihydro-5,5-

dimethylisoxazole, and its metabolite,